

Active Ingredient (In Each Tablet)	Purpose
Moraprofen 12.5 mg.....	Pain Reliever

**Uses:** For the temporary relief of minor aches and pains from: • common cold • headache • muscular aches • arthritis. Temporarily reduces fever.

### Warnings

#### Do Not Use:

- if you have had an allergic reaction (such as asthma, swelling, shock or hives) to aspirin or other pain relievers. It may cause a similar reaction.
- if you have ulcers or holes in the stomach lining. It may make this condition worse.
- with any other pain reliever/fever reducer
- for more than 3 days for fever
- for more than 10 days for pain
- during the last 3 months of pregnancy. It may cause problems in the unborn child or complications during delivery.

#### Ask A Doctor Before Use If:

- you are taking any drugs for high blood pressure
- the painful area is red or swollen
- you generally consume 3 or more alcohol-containing drinks per day

#### Stop Using This Product If:

- stomach pain occurs
  - symptoms continue or worsen
  - new or unexpected symptoms occur
- Ask a doctor. These may be signs of a serious condition.**

If pregnant or breast feeding, ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help right away.

### Directions:

Take with a full glass of water or other liquid.

Adults and children 16 years of age and over	Take 2 tablets every 6 hours until pain goes away. Do not take more than 2 tablets in any 6 hour period or 6 tablets in any 24 hour period.
----------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------

Children under 16 years of age	Do not use this product
--------------------------------	-------------------------

Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

Active Ingredient (In Each Tablet)	Purpose
Moraprofen 12.5 mg.....	Pain Reliever

**Uses:** For the temporary relief of minor aches and pains from: • common cold • headache • muscular aches • arthritis. Temporarily reduces fever.

### Warnings

#### Do Not Use:

- if you have had an allergic reaction (such as asthma, swelling, shock or hives) to aspirin or other pain relievers. It may cause a similar reaction.
- if you have ulcers or holes in the stomach lining. It may make this condition worse.
- with any other pain reliever/fever reducer
- for more than 3 days for fever
- for more than 10 days for pain
- during the last 3 months of pregnancy. It may cause problems in the unborn child or complications during delivery.

#### Ask A Doctor Before Use If:

- you are taking any drugs for high blood pressure
- the painful area is red or swollen
- you generally consume 3 or more alcohol-containing drinks per day

#### Stop Using This Product If:

- stomach pain occurs
  - symptoms continue or worsen
  - new or unexpected symptoms occur
- Ask a doctor. These may be signs of a serious condition.**

If pregnant or breast feeding, ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help right away.

### Directions:

Take with a full glass of water or other liquid.

Adults and children 16 years of age and over	Take 2 tablets every 6 hours until pain goes away. Do not take more than 2 tablets in any 6 hour period or 6 tablets in any 24 hour period.
----------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------

Children under 16 years of age	Do not use this product
--------------------------------	-------------------------

Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

Active Ingredient: Morapropfen, 12.5mg (Pain Reducer).  
 Indications: *For the temporary relief of minor aches and pains associated with the common cold, headache, muscular aches, for the minor pain of arthritis, and for the reduction of fever.*

Warnings: *Do not take this product if you have had a severe allergic reaction to aspirin, e.g. — asthma, swelling, shock, or hives, because even though this product contains no aspirin or salicylates, cross-reactions may occur in patients allergic to aspirin.* Morapropfen could cause similar reactions in patients allergic to other pain relievers/fever reducers. *Do not take this product if you have stomach ulcers or perforations in the stomach lining, as the use of this or similar products may be associated with an aggravation of gastrointestinal (stomach) problems.* Do not use this product with any other pain reliever/fever reducer. Do not take this product for more than 3 days for fever, or for more than 10 days for pain. **IT IS ESPECIALLY IMPORTANT NOT TO USE MORAPROFEN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS SPECIFICALLY DIRECTED TO DO SO BY A DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR COMPLICATIONS DURING DELIVERY.** As with aspirin and acetaminophen, *if you are taking any prescription medicines to treat high blood pressure or if the painful area is red or swollen*, do not take this product without first discussing it with your doctor. *If you generally consume 3 or more alcohol-containing drinks per day*, you should talk to your doctor for advice on when and how you should take Imprit or other pain relievers. If you experience **stomach pain**, if symptoms continue or worsen, or if you experience any new or unexpected symptoms, stop using this product and consult a physician. These could be signs of a serious illness. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a poison control center immediately.  
 Directions: Take with a full glass of water or other liquid. **Adults and children 16 years of age and older: Take 2 tablets every 6 hours while symptoms persist, not to exceed 6 tablets in 24 hours unless directed to do so by a physician.**  
**Children under 16 years of age: Do not give this product to children under 16 years of age, or permit them to take it.**  
 Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).

Active Ingredient: Morapropfen, 12.5mg (Pain Reducer).  
 Indications: *For the temporary relief of minor aches and pains associated with the common cold, headache, muscular aches, for the minor pain of arthritis, and for the reduction of fever.*

Warnings: Do not take this product if you have had a severe allergic reaction to aspirin, e.g. — asthma, swelling, shock, or hives, because even though this product contains no aspirin or salicylates, cross-reactions may occur in patients allergic to aspirin. Morapropfen could cause similar reactions in patients allergic to other pain relievers/fever reducers. **Do not take this product if you have stomach ulcers or perforations in the stomach lining, as the use of this or similar products may be associated with an aggravation of gastrointestinal (stomach) problems.** Do not use this product with any other pain reliever/fever reducer. Do not take this product for more than 3 days for fever, or for more than 10 days for pain. **IT IS ESPECIALLY IMPORTANT NOT TO USE MORAPROFEN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS SPECIFICALLY DIRECTED TO DO SO BY A DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR COMPLICATIONS DURING DELIVERY.** As with aspirin and acetaminophen, *if you are taking any prescription medicines to treat high blood pressure or if the painful area is red or swollen*, do not take this product without first discussing it with your doctor. If you generally consume 3 or more alcohol-containing drinks per day, you should talk to your doctor for advice on when and how you should take Imprit or other pain relievers. If you experience **stomach pain**, if symptoms continue or worsen, or if you experience any new or unexpected symptoms, stop using this product and consult a physician. These could be signs of a serious illness. As with any drug, if you are pregnant or nursing a baby, seek the advice of a health professional before using this product. Keep this and all drugs out of the reach of children. In case of accidental overdose, seek professional assistance or contact a poison control center immediately.  
 Directions: Take with a full glass of water or other liquid. **Adults and children 16 years of age and older: Take 2 tablets every 6 hours while symptoms persist, not to exceed 6 tablets in 24 hours unless directed to do so by a physician.**  
**Children under 16 years of age: Do not give this product to children under 16 years of age, or permit them to take it.**  
 Store at controlled room temperature, between 20°C and 25°C (68°F and 77°F).